

SUPPLEMENTAL MATERIAL

Supplementary Text

Cohort descriptions

Genotyping methods

Secondary Statistical Analyses

Acknowledgements

Supplemental Figures

Figure S1. Quantile-quantile plot of coronary artery calcification quantity single variant association.

Figure S2. Regional association plots of coronary artery calcification quantity at 9p21 (*CDKN2B*).

Figure S3. Regional association plots of coronary artery calcification quantity at 6p24 (*PHACTR1*).

Figure S4. Regional association plots of coronary artery calcification quantity at 2p24 (*APOB*).

Figure S5. Regional association plots of coronary artery calcification quantity at 19q13 (*APOE*).

Figure S6. Quantile-quantile plots of coronary artery calcification quantity gene-based association.

Figure S7. Quantile-quantile plot of carotid intima media thickness single variant association.

Figure S8. Regional association plots of carotid intima media thickness at 19q13 (*APOE*).

Figure S9. Quantile-quantile plot of carotid intima media thickness gene-based association.

Figure S10. Association of major *APOE* genotypes with CAC.

Supplemental Tables

Table S1. Demographics of Participants by Cohort in CAC Analysis.

Table S2. Demographics of Participants by Cohort in CIMT Analysis.

Table S3. *APOE* $\epsilon 2$ association with CAC adjusted for blood lipids.

Table S4. CHD-associated variants and association with subclinical atherosclerosis

Supplemental References

Cohort descriptions

Cardiovascular Health Study (CHS)

The CHS is a population-based cohort study of risk factors for CHD and stroke in adults ≥ 65 years conducted across four field centers in the United States. [The original predominantly European-ancestry cohort of 5201 persons was recruited in 1989-1990 from a random sample of people on Medicare eligibility lists and an additional 687 African-Americans were enrolled subsequently for a total sample of 5888. Blood samples were drawn from all participants at their baseline examination and DNA was subsequently extracted from available samples. Genotyping was performed at the General Clinical Research Center's Phenotyping/Genotyping Laboratory at Cedars-Sinai among CHS participants who consented to genetic testing and had DNA available using the Illumina HumanExome v.1.0 BeadChip array. Genotypes were jointly called at The University of Texas Health Science Center at Houston as part of approximately 62,000 ethnically-diverse samples from the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium.¹ Analyses were performed separately for individuals of European and African ancestries.

Diabetes Heart Study (DHS)

The Diabetes Heart Study (DHS) is a family-based observational cohort study of cardiovascular disease from a single research center in the United States.² The original predominantly (85%) European-ancestry cohort of 1443 persons was recruited in 1997-2005 from a random from families with at least two type 2 diabetes affected siblings and, if possible, a non-diabetic sibling. Blood samples were drawn from all participants at their baseline examination and DNA was subsequently extracted from available samples. Genotyping was performed on the Illumina HumanExome v.1.0 BeadChip array at the Center for Genomics and Personalized Medicine Research at Wake Forest University School of Medicine. Genotypes were called using ZCall³ and QC was performed at Wake Forest University Center for Genomics and Personalized Medicine Research.

Jackson Heart Study (JHS)

The JHS aimed at enrolling representative, population-based cohort of self-defined African-American persons aged 35–84 years, with an embedded collection of families for genetic study.⁴ Participants were enrolled from the three counties that make up the Jackson, Mississippi metropolitan area. Relatives of selected participants were recruited to develop a large, nested family cohort. Participants provided extensive medical and social history, had an array of physical and biochemical measurements and diagnostic procedures, and provided genomic DNA. Data and biologic materials have been collected from 5302 adult African Americans, including 1499 members of 291 families. Participants have a high prevalence of diabetes, hypertension, obesity, and related disorders. Genotyping was performed among JHS participants who consented to genetic testing and had DNA available using the Illumina HumanExome v.1.0 BeadChip array. Genotypes were jointly called at The University of Texas Health Science Center at Houston as part of approximately 62,000 ethnically-diverse samples from the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium.¹ Individuals that overlapped with ARIC were removed.

Family Heart Study (FamHS)

The FamHS (<https://dsgweb.wustl.edu/fhsc/>) began in 1992 with the ascertainment of 1,200 families, half randomly sampled and half selected because of an excess of CHD or risk factor abnormalities as compared with age- and sex-specific population rates.⁵ The families, with approximately 6,000 subjects, were sampled from four population-based parent studies: the Framingham Heart Study, the Utah Family Tree Study, and two centers for the Atherosclerosis Risk in Communities study (ARIC: Minneapolis, and Forsyth County, NC). These individuals attended a clinic visit between the years 1994-1996 and a broad range of phenotypes were assessed in the general domains of CHD, atherosclerosis, cardiac and vascular function, inflammation and hemostasis, lipids and lipoproteins, blood pressure, diabetes and insulin resistance, pulmonary function, diet, habitual physical activity, anthropometry, medical history and medication use. Approximately 8 years later, study participants belonging to the largest pedigrees were invited for a second clinical exam (2002-04) and computed tomography scan measurements were assessed, among many other phenotypes. A total of 2,756 European American (EA) subjects in 510 extended random and high CHD risk families were studied. A total of 1,865 key EA subjects within this group of families were genotyped using the Illumina Infinium HumanExome v1.0 BeadChip. In addition, 633 African American (AA) subjects were recruited at an additional ARIC field center at the University of Alabama in Birmingham. A total of 608 AA subjects were genotyped using the Illumina Infinium HumanExome v1.0 BeadChip. Informed consent was obtained from all participants, and this project was approved by the Institutional Review Boards of all participating institutions.

Lothian Birth Cohort 1936 (LBC1936)

The LBC1936 consists of 1,091 relatively healthy individuals assessed on cognitive and medical traits at about 70 years of age. They were all born in 1936 and most took part in the Scottish Mental Survey of 1947. At baseline the sample of 548 men and 543 women had a mean age 69.6 years (SD = 0.8). They were all Caucasian, community-dwelling, and almost all lived in the Lothian region (Edinburgh city and surrounding area) of Scotland. A full description of participant recruitment and testing can be found elsewhere.⁶ 988 individuals were genotyped using the Illumina HumanExome BeadChip at the Wellcome Trust Clinical Research Facility, Edinburgh.

SHIP / SHIP-Trend

SHIP is a population-based project in West Pomerania, a region in the northeast of Germany, that consists of two independent prospectively collected cohorts (SHIP and SHIP-TREND) assessing the prevalence and incidence of common population-based diseases and their risk factors. The study design has been previously described in detail.⁷ Briefly, a sample from the population aged 20 to 79 years was drawn from population registries. First, the three cities of the region (with 17,076 to 65,977 inhabitants) and the 12 towns (with 1,516 to 3,044 inhabitants) were selected, and then 17 out of 97 smaller towns (with less than 1,500 inhabitants), were drawn at random. Second, from each of the selected communities, subjects were drawn at random, proportional to the population size of each community and stratified by age and gender. Only individuals with German citizenship and main residency in the study area were included. For SHIP, baseline examinations were carried out from 1997 until 2001, and the sample finally comprised 4,308 participants. Baseline examinations for SHIP-TREND were carried out between 2008 and 2012, finally comprising 4420 participants. Analyses were performed separately for individuals of both cohorts.

Genetic Study of Atherosclerosis Risk (GeneSTAR)

GeneSTAR is a family-based prospective study of risk factors, occult disease, and incident cardiovascular disease in siblings, later extended to offspring and whole pedigrees. European- and African American families were identified from probands with early-onset (< age 60) coronary artery disease (CAD) hospitalized in any of 10 Baltimore hospitals between 1983 and 2007. Siblings completed baseline screening between 1983 and 2007, while offspring, co-parents of the offspring, and additional siblings completed baseline screening between 2003 and 2007. Siblings were followed every 5 years for cardiovascular and other comorbid incident events. DNA was collected at baseline (1991-2007) or follow-up (for those whose baseline visit was pre-1991). Probands were not eligible if they had CAD associated with calcific aortic stenosis or chronic glucocorticosteroid therapy, following organ transplantation or post intensive chest radiation, had an autoimmune diseases like systemic lupus, or if they had a cocaine-induced myocardial infarction. Participants younger than age 21 or older than age 80, who had known CAD or an autoimmune disease such as systemic lupus, were taking chronic glucocorticosteroids, had undergone any organ transplantation, or had major comorbidity that had a life expectancy under 5 years of age were excluded. The full sample includes 4423 participants, 51% female/49% male, and 38% African American/61% European American/1% other American. Genotyping was performed by the Northwest Genomics Center at the University of Washington through the RS&G service using the Illumina HumanExome v.1.2 BeadChip array. Analyses were performed separately for European and African Americans.

ImaGene

Project ‘Cardiovascular phenotype-genotype analysis with a CT based lung cancer screening trial’ within the Population Imaging Genetics (ImaGene) study was set up to investigate the relationship between genetics and image characteristics related to cardiovascular disease in an at-risk population of current or former heavy smokers between 50 and 75 years of age. 905 participants of the Dutch-Belgian lung cancer screening trial (NELSON) were successfully genotyped using the Illumina HumanExome v1.1 BeadChip array. Low-dose, non-ECG synchronized, non-contrast enhanced baseline chest CTs from the NELSON study were available for all participants. CTs were acquired on a 16 detector-row scanner (Mx8000 IDT, Somatom Sensation 16, or Brilliance 16P, Philips Medical Systems, Cleveland, OH, USA) in spiral mode with 16×0.75 mm collimation. Axial images of 1.0 mm thickness at 0.7 mm increment were reconstructed with a moderately soft kernel (Philips “B”). The peak voltage was 120–140 kVp depending on patient weight, with a tube current of 30 mAs.

Old Order Amish

The Amish Family Calcification Study (AFCS) was initiated in 2001 to identify the determinants of vascular calcification and to evaluate the relationship between calcification of bone and vascular tissue among members of the Old Order Amish community in Lancaster County.⁸ For this study relatively healthy subjects and their family members were recruited from the Amish community between 2001 and 2006 for vascular imaging by electron beam CT scan. The final sample size comprised 1,075 subjects. Blood samples were drawn for biochemical assays and DNA analysis.

Framingham Heart Study (FHS)

Framingham Heart Study (FHS) is a community-based prospective study designed to investigate the incidence of cardiovascular diseases and factors related to its development. Study population was composed of three cohorts with European ancestry, original (5,209 participants ascertained systematically from two-thirds of the households in the town of Framingham, MA, beginning in 1948), Offspring (5,124 children of the original cohort, and spouses of those children, beginning in 1972), and the Third Generation (4,095 children of the Offspring cohort, beginning in 2002).⁹
¹⁰ In addition to the initial examination and face-to-face interview, participants were invited to follow-up examinations at the study clinics that were accomplished every two years for the original cohort or every four years for the Offspring and Gen3 cohorts. An un-related spouse cohort was formed in 2004 include 103 spouses of the offspring participants. Genotyping for more than 276,000 variants on the basis of the Illumina Infinium Human Exome Array v1.0 or v1.1 was conducted in 8351 FHS samples. Quality control of the genotype was accomplished through jointly called at The University of Texas Health Science Center at Houston as part of approximately 62,000 ethnically-diverse samples from the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Consortium.¹

BioImage – High Risk Plaque

The BioImage study (NCT00738725) is a multi-ethnic, observational study aimed at characterizing subclinical atherosclerosis in 6,699 US adults (55-80 years at baseline, 2008-2009) at risk for, but without, clinical ASCVD.¹¹ Participants were genotyped on the Illumina Infinium Human Exome Array v1.1.¹²

Multi-Ethnic Study of Atherosclerosis (MESA)

MESA is an NHLBI-sponsored population-based, prospective, multi-center cohort study including participants recruited from six field sites in the United States – Forsyth County, NC (Wake Forest University), Northern Manhattan/Bronx, NY (Columbia University), Baltimore/Baltimore County, MD (Johns Hopkins University), St. Paul, MN (University of Minnesota, Twin Cities), Chicago, IL (Northwestern University), and Los Angeles County, CA (UCLA). Details of recruitment and study design have been previously published elsewhere.¹³ Briefly, the MESA cohort comprises 6,814 men and women of diverse ethnic background who were 45 to 84 years old at the baseline exam and free of clinically overt cardiovascular disease (CVD) who were recruited to elucidate the determinants and natural history of subclinical CVD, study progression of subclinical CVD, and its impact on incident clinical CVD. The cohort is 53% women with an ethnic composition of approximately 38% Caucasian, 28% African American, 22% Hispanic and 12% Asian, primarily of Chinese descent. Five clinical exams have taken place (2000-02, 2002-04, 2004-05, 2005-07, 2010-12), with follow-up every 9 to 12 months for events. The MESA was approved by the Institutional Review Board of all participating field sites and reading centers and participants gave informed consent for participation and use of DNA specimens. The current study includes data from African-Americans and Caucasians with available exome chip data, coronary artery calcium (CAC), and carotid intima media thickness (IMT) at the baseline exam (2000-02). Joint genotype calling for the exome chip V1.0 was performed at UT-Houston for all CHARGE samples, including for MESA. Additional QC was performed at Cedars-Sinai and University of Virginia. After QC, 238,895 SNPs remained for analysis.

Genetic Epidemiology Network of Arteriopathy (GENOA)

GENOA is one of four networks in the NHLBI Family-Blood Pressure Program (FBPP).¹⁴ GENOA's long-term objective is to elucidate the genetics of target organ complications of hypertension, including both atherosclerotic and arteriolosclerotic complications involving the heart, brain, kidneys, and peripheral arteries. The longitudinal GENOA Study recruited European-American and African-American sibships with at least 2 individuals with clinically diagnosed essential hypertension before age 60 years. All other members of the sibship were invited to participate regardless of their hypertension status. Participants were diagnosed with hypertension if they had either 1) a previous clinical diagnosis of hypertension by a physician with current anti-hypertensive treatment, or 2) an average systolic blood pressure ≥ 140 mm Hg or diastolic blood pressure ≥ 90 mm Hg based on the second and third readings at the time of their clinic visit. Exclusion criteria were secondary hypertension, alcoholism or drug abuse, pregnancy, insulin-dependent diabetes mellitus, or active malignancy. During the first exam (1995-2000), 1,583 European Americans from Rochester, MN and 1,854 African Americans from Jackson, MS were examined. Between 2000 and 2005, 1,241 of the European Americans returned for an examination that included a CT scan of the heart for coronary artery calcification. Between 2009 and 2011, 657 of the African Americans returned for an examination that included a CT scan of the heart for coronary artery calcification. Because African-American probands for GENOA were recruited through the Atherosclerosis Risk in Communities (ARIC) Jackson field center participants, we excluded ARIC participants from analyses. Written informed consent was obtained from all participants, and this project was approved by the Institutional Review Boards of all participating institutions. Both European Americans and African Americans were genotyped using the Illumina Infinium HumanExome v1.0 BeadChip. Genotyping and genotype calling for European Americans was performed at the University of Texas Health Sciences Center. Genotyping and genotype calling for African Americans was performed at the Center for Inherited Disease Research (CIDR).

AGES Reykjavik

The Reykjavik Study cohort originally comprised a random sample of 30,795 men and women born in 1907–1935 and living in Reykjavik in 1967. A total of 19381 attended, resulting in 71% recruitment rate. The study sample was divided into six groups by birth year and birth date within month. One group was designated for longitudinal follow-up and was examined in all stages. One group was designated a control group and was not included in examinations until 1991. Other groups were invited to participate in specific stages of the study. Between 2002 and 2006, the AGES-Reykjavik study re-examined 5764 survivors of the original cohort who had participated before in the Reykjavik Study. The AGES Reykjavik Study GWAS was approved by the National Bioethics Committee (00-063- V8+1) and the Data Protection Authority. DNA was genotyped using the Illumina HumanExome v.1.0 BeadChip array. Samples were excluded from the dataset based on sample failure, genotype mismatch with reference panel, and sex mismatch, resulting in clean genotype data on 2,983 individuals. Standard protocols for working with Illumina data were followed, with clustering score greater than 0.4. SNPs were excluded using filters based on call rate ($<97\%$), Hardy-Weinberg Equilibrium ($<1 \times 10^{-6}$) 238,015 SNPs passing all QC (of 247,039 prior to cleaning steps).

Atherosclerosis Risk in Communities (ARIC)

ARIC is a population-based prospective study of men and women aged 45-64 years at baseline, which recruited 15,792 African American and White individuals from 4 U.S. communities to study the etiology and natural history of subclinical and clinical atherosclerosis. Participants

were examined at baseline (1987-1989) and 4 follow-up clinic visits (1990-1992, 1993-1995, 1996-1998, 2011-2013), at which a rich array of health assessments have been made, with over 25 years of follow-up through annual phone updates for hospital admissions, medical history, and events.¹⁵ DNA has been collected, with extensive genome-wide array and genetic sequencing data available.

Rotterdam Study (RS)

The Rotterdam Study (RS) is a prospective population-based cohort study among persons aged 55 years or older in the Ommoord district of Rotterdam, the Netherlands. The rationale and design of the Rotterdam Study have been described elsewhere.¹⁶ The cohort started in 1990 and has been extended twice. The present study used data for carotid intima media thickness from the baseline examination of the original cohort (RS-I, visit 1: 1990-1993) and for coronary calcification from the third examination of the original cohort (RS-I, visit 3: 1997-1999). The RS participants are interviewed and have an extensive set of examinations every 3-4 years and have been followed-up for a variety of diseases. The Rotterdam Study has been approved by the Medical Ethics Committee of the Erasmus MC and by the Ministry of Health, Welfare and Sport of the Netherlands, implementing the Wet Bevolkingsonderzoek: ERGO (Population Studies Act: Rotterdam Study). All participants provided written informed consent to participate in the study and to obtain information from their treating physicians. The approval has been renewed every 5 years, as well as with the introduction of major new elements in the study. The methods regarding the Exome Chip Array has been described in detail elsewhere.¹

Erasmus Rucphen Family Study (ERF)

Erasmus Rucphen Family study (ERF) is a family based study conducted in a genetically isolated population in the South-West of the Netherlands, studied as part of the Genetic Research in Isolated Population (GRIP) program.^{17, 18} The aim of this study is to identify genetic risk factors of complex diseases and genetic associations to complex traits. Study population includes approximately 3,000 participants who are descendants of a limited number of founders living in the 19th century. All data were collected between 2002 and 2005. All participants gave written informed consent and the Medical Ethics Committee at Erasmus MC University Medical Center approved the study. Study participants from the ERF cohort (N = 1,527) were genotyped on the Illumina Infinium HumanExome BeadChip, version 1.1. Calling was performed with GenomeStudio and the ZCall variant calling tool (Broad Institute).³

Netherlands Epidemiology of Obesity Study (NEO)

The NEO was designed for extensive phenotyping to investigate pathways that lead to obesity-related diseases. The NEO study is a population-based, prospective cohort study that includes 6,671 individuals aged between 45-65 years, with an oversampling of individuals with overweight or obesity. At baseline, information on demography, lifestyle, and medical history have been collected by questionnaires. In addition, samples of 24-h urine, fasting and postprandial blood plasma and serum, and DNA were collected. Genotyping was performed using the Illumina HumanCoreExome chip, which was subsequently imputed to the 1000 genome reference panel. Participants underwent an extensive physical examination, including anthropometry, electrocardiography, spirometry, and measurement of the carotid artery intima-media thickness by ultrasonography. In random subsamples of participants, magnetic resonance imaging of abdominal fat, pulse wave velocity of the aorta, heart, and brain, magnetic resonance

spectroscopy of the liver, indirect calorimetry, dual energy X-ray absorptiometry, or accelerometry measurements were performed. The collection of data started in September 2008 and completed at the end of September 2012. Participants were currently being followed for the incidence of obesity-related diseases and mortality.¹⁹

Young Finns Study (YFS)

The Cardiovascular Risk in Young Finns Study (YFS) is a population-based follow up-study started in 1980. The main aim of the YFS is to determine the contribution made by childhood lifestyle, biological and psychological measures to the risk of cardiovascular diseases in adulthood. In 1980, over 3,500 children and adolescents all around Finland participated in the baseline study. The follow-up studies have been conducted mainly with 3-year intervals. The latest 30-year follow-up study was conducted in 2010-11 (ages 33-49 years) with 2,063 participants. The study was approved by the local ethics committees (University Hospitals of Helsinki, Turku, Tampere, Kuopio and Oulu) and was conducted following the guidelines of the Declaration of Helsinki. All participants gave their written informed consent.²⁰

Genotyping methods

Genotypes from the Age, Gene/Environment, Susceptibility-Reykjavik (AGES) Study, Cardiovascular Health Study (CHS), Rotterdam Study (RS), Framingham Heart Study (FHS), Family Heart Study (FamHS), and Jackson Heart Study (JHS) were jointly called.¹ High Risk Plaque-Bioimage (Bioimage), genotyped with v1.1 of the Beadchip, was called in GenomeStudio (Illumina) using GenCall and low frequency variant calling was further refined using zCall.³ The Diabetes Heart Study (DHS), Genetic Epidemiology Network of Arteriopathy Study (GENOA), ImaGene, Young Finns Study (YFS), Gene Study of Atherosclerosis Risk in Families (GeneSTAR), Old Order Amish Study (Amish), Netherlands Epidemiology of Obesity Study (NEO), Erasmus Ruchphen Family Study, Lothian Birth Cohort, Study of Health in Pomerania (SHIP), and SHIP-TREND were called in GenomeStudio (Illumina) using the joint calling cluster file and used zCall.

It is estimated that 95 % of European ancestry and 85% of African ancestry variation with minor allele frequency (MAF) > 0.1 % is captured by the content of the Illumina HumanExome Beadchip array.²¹

Variant quality control was performed jointly¹ and by individual cohorts. Low-performing variants from joint-calling were excluded from analysis.¹ 238,065 variants were included in analysis. Samples were excluded by each cohort if genotypes had poor concordance with prior genome-wide array-based genotypes when available, >5% of genotypes missing, they were population clustering outliers, had high inbreeding coefficients or heterozygosity rates, phenotypic and genotypic sex discordance, one from each duplicate (or monozygotic twin pairing), and high degrees of cryptic relatedness in studies without families was observed.²¹ All variants were coded in an additive fashion. Variants were annotated using dbNSFP v2.0.²²

Secondary Statistical Analyses

***APOE* ϵ 2 Association Conditional on LDL cholesterol**

To determine whether low-density lipoprotein (LDL) cholesterol is an important intermediary in the association between rs7412-T and CAC, we tested the association further adjusting for LDL cholesterol. In studies with available measures, fasting LDL cholesterol was obtained and/or calculated based on the Friedewald calculation when triglycerides were less than 400 mg/dL and directly measured when triglycerides were greater than 400 mg/dL. To account for the effect of lipid-lowering therapy, we adjusted LDL cholesterol to reflect the observation that statins, on average, reduce LDL cholesterol by 30% prior to conditioning.²³ Association analyses were performed within each cohort and meta-analyzed as described above with fasting LDL cholesterol as an additional covariate.

***APOE* ϵ 2 Association Conditional on Other *APOE* Genotypes**

The major *APOE* polymorphisms at 19q13.2 are ϵ 2, ϵ 3, and ϵ 4.²⁴ The ϵ 2 allele is determined by rs7412-T and the ϵ 4 allele by rs429358-C, while the referent ϵ 3 genotype has the rs7412-C / rs429358-T haplotype. Given the absence of rs429358 on the exome chip and prior genome-wide genotyping arrays, we obtained directly genotyped ϵ 2, ϵ 3, and ϵ 4 by polymerase chain reaction in 5,872 participants from FHS, AGES, GENOA, CHS, and DHS. Our hypothesis was that the effect of the ϵ 2 allele on subclinical atherosclerosis was independent of ϵ 4 status. We tested the association of the *APOE* non-reference genotypes (ϵ 3/ ϵ 3 referent) with CAC stratified for ethnicity and sex while accounting for age and principal components within each cohort. We then performed a fixed-effects meta-analysis weighted by the inverse of variances to summarize the association across cohorts.

Association with CHD

To test for association of rs7412-T with prevalent CHD, we obtained summary results from an association analysis led by the Myocardial Infarction Genetics (MIGen) Consortium. 21,182 individuals of European ancestry, independent of the subclinical atherosclerosis analyses, were genotyped with the Illumina HumanExome Beadchip v1.0 at the Broad Institute (Cambridge, MA) and 9,472 of these individuals were classified as having CHD.²⁵ We obtained the single variant association results for rs7412-T accounting for age, sex, ethnicity, and principal components of ancestry.

We also determined the concordance of subclinical atherosclerosis genetic architecture with that of CHD. We focused on the concordance of effect direction of variants previously associated with CHD at a genome-wide level for individuals of European ancestry.²⁶ We selected the top associated CHD variant for each locus genotyped on the exome chip. Separately for CAC and CIMT, we determined the probability of a CHD variant having a concordant effect direction with each subclinical atherosclerotic trait. We determined the likelihood of effect observed concordance compared to chance accounting for each variant's risk allele's frequency with an exact binomial test.

Acknowledgements / Funding

Cardiovascular Health Study (CHS)

This Cardiovascular Health Study (CHS) research was supported by NHLBI contracts HHSN268201200036C, HHSN268200800007C, N01HC55222, N01HC85079, N01HC85080, N01HC85081, N01HC85082, N01HC85083, N01HC85086, N01HC85085, N01HC45133; and NHLBI grants U01HL080295, R01HL087652, R01HL105756, R01HL103612, R01HL120393, R01HL130114, and R01HL068986 with additional contribution from the National Institute of Neurological Disorders and Stroke (NINDS). Additional support was provided through R01AG023629 from the National Institute on Aging (NIA). A full list of principal CHS investigators and institutions can be found at CHS-NHLBI.org. The provision of genotyping data was supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR000124, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Diabetes Heart Study (DHS)

The Diabetes Heart Study (DHS) research reported in this article was supported in part by R01 HL67348, R01 HL092301, R01 NS058700 and the General Clinical Research Centre of the Wake Forest School of Medicine (M01 RR07122, F32 HL085989). The authors thank the other investigators, the staff, and the participants of the DHS study for their valuable contributions.

Jackson Heart Study (JHS)

We thank the Jackson Heart Study (JHS) participants and staff for their contributions to this work. The JHS is supported by contracts HHSN268201300046C, HHSN268201300047C, HHSN268201300048C, HHSN268201300049C, HHSN268201300050C from the National Heart, Lung, and Blood Institute and the National Institute on Minority Health and Health Disparities.

Family Heart Study (FamHS)

This Family Heart Study (FamHS) research was supported by NIH grants R01-HL-117078, R01-HL-087700 and R01-HL-088215 from NHLBI; and R01-DK-089256 from NIDDK.

Lothian Birth Cohort 1936 (LBC1936)

We thank the cohort participants and team members who contributed to this study. Genotyping was supported by Centre for Cognitive Ageing and Cognitive Epidemiology (Pilot Fund award), Age UK and the Royal Society of Edinburgh and was conducted at the Wellcome Trust Clinical Research Facility Genetics Core at Western General Hospital, Edinburgh, UK. Phenotype collection was supported by Age UK (The Disconnected Mind project). The work was undertaken by The University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology, part of the cross council Lifelong Health and Wellbeing Initiative (MR/K026992/1). Funding from the BBSRC and MRC is gratefully acknowledged.

SHIP / SHIP-Trend

SHIP is part of the Community Medicine Research net of the University of Greifswald, Germany, which is funded by the Federal Ministry of Education and Research (grants no. 01ZZ9603, 01ZZ0103, and 01ZZ0403), the Ministry of Cultural Affairs as well as the Social Ministry of the Federal State of Mecklenburg-West Pomerania, and the network ‘Greifswald Approach to Individualized Medicine (GANI_MED)’ funded by the Federal Ministry of Education and Research (grant 03IS2061A). ExomeChip data have been supported by the Federal Ministry of Education and Research (grant no. 03Z1CN22) and the Federal State of Mecklenburg-West Pomerania. The University of Greifswald is a member of the Caché Campus program of the InterSystems GmbH.

Genetic Study of Atherosclerosis Risk (GeneSTAR)

GeneSTAR was supported by grants from the National Institutes of Health/National Heart, Lung, and Blood Institute (HL112064, U01 HL72518, HL097698, HL59684, HL071025, HL092165, HL099747, and K23HL105897) and by a grant from the National Institutes of Health/National Center for Research Resources (M01-RR000052) to the Johns Hopkins General Clinical Research Center. Genotyping services were provided through the RS&G Service by the Northwest Genomics Center at the University of Washington, Department of Genome Sciences, under U.S. Federal Government contract number HHSN268201100037C from the National Heart, Lung, and Blood Institute.

ImaGene

The ImaGene study was financially supported by the Netherlands Organization for Scientific Research (NWO)/ Foundation for Technological Sciences (STW); Project 12726. CT scans were acquired within the NELSON study.

Old Order Amish

Supported by NIH grants: R01 HL69313, R01 HL088119, and P30 DK072488.

Framingham Heart Study (FHS)

The National Heart, Lung and Blood Institute’s Framingham Heart Study is supported by contract N01-HC-25195

BioImage – High Risk Plaque

The High Risk Plaque (HRP) Initiative encompassing the BioImage Study is a precompetitive industry collaboration funded by Abbott, Abbvie, AstraZeneca, BG Medicine, Merck, Philips, and Takeda. HRP Joint Steering Committee: Pieter Muntendam, MD (BG Medicine); Aram Adourian (BG Medicine); Michael Klimas, PhD (Merck); Joel Raichlen, MD (AstraZeneca); Oliver Steinbach (Philips); James Beckett (Philips); Ramon Espaillet (Abbvie); Michael Jarvis (Abbvie) and Tomoyuki Nishimoto (Takeda). The sponsor had no role in the study design; in the collection, analysis, and interpretation of the data; in the writing of this report; or in the decision to submit the paper for publication.

Multi-Ethnic Study of Atherosclerosis (MESA)

MESA and the MESA SHARe project are conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts HHSN268201500003I, N01-HC-95159, N01-HC-95160, N01-HC-95161,

N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-001079, UL1-TR-000040, and DK063491. MESA Family is conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support is provided by grants and contracts R01HL071051, R01HL071205, R01HL071250, R01HL071251, R01HL071258, R01HL071259, by the National Center for Research Resources, Grant UL1RR033176, and the National Center for Advancing Translational Sciences, Grant UL1TR000124. Funding for SHARe genotyping was provided by NHLBI Contract N02-HL-64278.

Genetic Epidemiology Network of Arteriopathy (GENOA)

Support for GENOA was provided by the National Heart, Lung and Blood Institute (HL119443, HL118305, HL054464, HL054457, HL054481, HL071917, HL085571, HL086694, and HL87660) of the National Institutes of Health. Genotyping was performed at the University of Texas Health Sciences Center (Eric Boerwinkle, Megan Grove-Gaona) and the Center for Inherited Disease Research (CIDR). We would like to thank the families that participated in the GENOA study.

AGES Reykjavik

This study has been funded by NIH contracts N01-AG-1-2100 and 271201200022C, the NIA Intramural Research Program, Hjartavernd (the Icelandic Heart Association), and the Althingi (the Icelandic Parliament). The study is approved by the Icelandic National Bioethics Committee, VSN: 00-063. The researchers are indebted to the participants for their willingness to participate in the study.

Atherosclerosis Risk in Communities (ARIC)

The Atherosclerosis Risk in Communities Study is carried out as a collaborative study supported by National Heart, Lung, and Blood Institute contracts (HHSN268201100005C, HHSN268201100006C, HHSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C), R01HL087641, R01HL59367 and R01HL086694; National Human Genome Research Institute contract U01HG004402; and National Institutes of Health contract HHSN268200625226C.

Rotterdam Study (RS)

The generation and management of the Illumina exome chip v1.0 array data for the Rotterdam Study (RS-I) was executed by the Human Genotyping Facility of the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, Rotterdam, The Netherlands. The Exome chip array data set was funded by the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, from the Netherlands Genomics Initiative (NGI)/Netherlands Organisation for Scientific Research (NWO)-sponsored Netherlands Consortium for Healthy Aging (NCHA; project nr. 050-060-810); the Netherlands Organization for Scientific Research (NWO; project number 184021007) and by the Rainbow Project (RP10; Netherlands Exome Chip Project) of the Biobanking and Biomolecular Research Infrastructure Netherlands (BBMRI-NL; www.bbmri.nl). We thank Ms. Mila Jhamai, Ms. Sarah Higgins, and Mr. Marijn Verkerk for their help in creating the exome chip database, and Carolina Medina-Gomez, MSc, Lennard Karsten, MSc, and Linda Broer PhD for QC and variant calling. Variants were called using the

best practice protocol developed by Grove et al. as part of the CHARGE consortium exome chip central calling effort [PMID:23874508]. The Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, Rotterdam, Netherlands Organization for the Health Research and Development (ZonMw), the Research Institute for Diseases in the Elderly (RIDE), the Ministry of Education, Culture and Science, the Ministry for Health, Welfare and Sports, the European Commission (DG XII), and the Municipality of Rotterdam. The authors are grateful to the study participants, the staff from the Rotterdam Study and the participating general practitioners and pharmacists.

Erasmus Rucphen Family Study (ERF)

The ERF study as a part of EUROSPAN (European Special Populations Research Network) was supported by European Commission FP6 STRP grant number 018947 (LSHG-CT-2006-01947) and also received funding from the European Community's Seventh Framework Programme (FP7/2007-2013)/grant agreement HEALTH-F4-2007-201413 by the European Commission under the programme "Quality of Life and Management of the Living Resources" of 5th Framework Programme (no. QLG2-CT-2002-01254). The ERF study was further supported by ENGAGE consortium and CMSB. High-throughput analysis of the ERF data was supported by joint grant from Netherlands Organisation for Scientific Research and the Russian Foundation for Basic Research (NWO-RFBR 047.017.043). Exome sequencing in ERF was supported by the ZonMw grant (project 91111025). Exome-chip genotyping was supported by BBMRI-NL, a Research Infrastructure financed by the Dutch Government (NWO 184.021.007). We are grateful to all study participants and their relatives, general practitioners and neurologists for their contributions and to P. Veraart for her help in genealogy, J. Vergeer for the supervision of the laboratory work and P. Snijders for his help in data collection.


Netherlands Epidemiology of Obesity Study (NEO)

The authors of the NEO study thank all individuals who participated in the Netherlands Epidemiology of Obesity study, all participating general practitioners for inviting eligible participants and all research nurses for collection of the data. We thank the NEO study group, Pat van Beelen, Petra Noordijk and Ingeborg de Jonge for the coordination, lab and data management of the NEO study. The genotyping in the NEO study was supported by the Centre National de Génotypage (Paris, France), headed by Jean-Francois Deleuze. The NEO study is supported by the participating Departments, the Division and the Board of Directors of the Leiden University Medical Center, and by the Leiden University, Research Profile Area Vascular and Regenerative Medicine. Dennis O. Mook-Kanamori is supported by Dutch Science Organization (ZonMW-VENI Grant 916.14.023)

Young Finns Study (YFS)


The Young Finns Study has been financially supported by the Academy of Finland: grants 286284, 134309 (Eye), 126925, 121584, 124282, 129378 (Salve), 117787 (Gendi), and 41071 (Skidi); the Social Insurance Institution of Finland; Kuopio, Tampere and Turku University Hospital Medical Funds (grant X51001); Juho Vainio Foundation; Paavo Nurmi Foundation; Finnish Foundation of Cardiovascular Research ; Finnish Cultural Foundation; Tampere Tuberculosis Foundation ; Emil Aaltonen Foundation ; and Yrjö Jahnsson Foundation. The expert technical assistance in the statistical analyses by Ville Aalto and Irina Lisinen is gratefully acknowledged.

Figure S1. Quantile-quantile plot of coronary artery calcification quantity.




The expected association p-values for the distribution of genotyped versus the observed distribution of p-values for CAC quantity association is displayed. Significant systemic inflation is not observed ($\lambda_{GC} = 1.057$). The dotted red line represents the pre-specified threshold for statistical significance accounting for the number of individual variants tested for association ($0.05 / 238,065 = 2.10 \times 10^{-7}$). Lead variants of novel significant associated loci (*APOE* p.R176C and *APOB* p.R3527Q) are highlighted in red.

A.




B.




The *CDKN2B* gene resides in the 9p21 genomic region. These regional association plots demonstrate the strength of association, by $-\log_{10}(\text{p-value})$, for the lead 9p21 variant, rs10757278, and each genotyped variant within +/- 400 kilobases. Color-coding based on ethnic-specific correlation observed in the 1000 Genomes Project is demonstrated. Dot sizes are scaled to the number genotyped for a given variant. Plots are separately shown for participants of: **A.** European ancestry and **B.** African ancestry.

Figure S3. Regional association plots of coronary artery calcification quantity at 6p24.

A.




B.




The *PHACTR1* gene resides in the 6p24 genomic region. These regional association plots demonstrate the strength of association, by $-\log_{10}(\text{p-value})$, for the lead 6p24 variant, rs9349379, and other genotyped variants within ± 400 kilobases. Color-coding based on ethnic-specific correlation observed in the 1000 Genomes Project is demonstrated. Dot sizes are scaled to the number genotyped for a given variant. Plots are separately shown for participants of: **A.** European ancestry and **B.** African ancestry.

Figure S4. Regional association plots of coronary artery calcification quantity at 2p24.

A.



B.



The *APOB* gene resides in the 2p24 genomic region. These regional association plots demonstrate the strength of association, by $-\log_{10}(\text{p-value})$, for the lead 2p24 variant, rs5742904, and each genotyped variant within +/- 400 kilobases. Color-coding based on ethnic-specific correlation observed in the 1000 Genomes Project is demonstrated. Dot sizes are scaled to the number genotyped for a given variant. Plots are separately shown for participants of: **A.** European ancestry and **B.** African ancestry.


A.




The *APOE* gene resides in the 19q13 genomic region. These regional association plots demonstrate the strength of association, by $-\log_{10}(\text{p-value})$, for the lead 19q13 variant, rs7412, and each genotyped variant within +/- 400 kilobases. Color-coding based on ethnic-specific correlation observed in the 1000 Genomes Project is demonstrated. Dot sizes are scaled to the number genotyped for a given variant. Plots are separately shown for participants of: **A.** European ancestry and **B.** African ancestry.

Figure S6. Quantile-quantile plots of coronary artery calcification quantity gene-based association.

A.




B.



The expected associations p-values for the distribution of genotyped versus the observed distributions of p-values for CAC quantity gene-based association is displayed. Systemic inflation is not observed (T1 $\lambda_{GC} = 1.026$, SKAT $\lambda_{GC} = 1.107$). Significant observed deviations from the expected distribution are not noted. Association analyses are displayed for: **A.** T1 and **B.** SKAT.


Figure S7. Quantile-quantile plot of carotid intima media thickness single variant association.




The expected association p-values for the distribution of genotyped versus the observed distribution of p-values for CIMT association is displayed. Systemic inflation is not observed ($\lambda_{GC} = 1.080$). The dotted red line represents the prespecified threshold for statistical significance accounting for the number of individual variants tested for association ($0.05 / 238,065 = 2.1 \times 10^{-7}$). The lead variant of the novel significant associated locus (*APOE* p.R176C) is highlighted in red.

Figure S8. Regional association plots of carotid intima media thickness at 19q13 (*APOE*).

A.




B.




The *APOE* gene resides in the 19q13 genomic region. These regional association plots demonstrate the strength of association, by $-\log_{10}(\text{p-value})$, for the lead 19q13 variant, rs7412, and each genotyped variant within +/- 400 kilobases. Color-coding based on ethnic-specific correlation observed in the 1000 Genomes Project is demonstrated. Dot sizes are scaled to the number genotyped for a given variant. Plots are separately shown for participants of: **A.** European ancestry and **B.** African ancestry.

Figure S9. Quantile-quantile plot of carotid intima media thickness gene-based association.
A.




B.



The expected associations p-values for the distribution of genotyped versus the observed distributions of p-values for CIMT gene-based association is displayed. Significant systemic inflation is not observed (T1 $\lambda_{GC} = 1.043$, SKAT $\lambda_{GC} = 1.114$). Significant observed deviations from the expected distribution are not noted. Association analyses are displayed for: **A.** T1 and **B.** SKAT.

Figure S10. Association of major *APOE* genotypes with CAC.



Major *APOE* genotypes ascertained by PCR were obtained in 5,872 individuals and were associated with log-transformed CAC quantity relative to the ε3/ε3 referent genotype within six cohorts accounting for age, sex, and ethnicity. Effect estimates were combined using fixed-effects meta-analysis and are displayed.

Table S1. Demographics of Participants by Cohort in CAC Analysis

Cohort	Ethnicity	Years measured	N	Age, y mean (sd)	Women, n (%)	CAC Score, median (IQR)	CAC>0, n (%)	CAC>100, n (%)	Hypertension, n (%)	Hypercholesterolemia, n (%)	Diabetes mellitus, n (%)	Recent cigarette smoker, n (%)	Device	References	
Cardiovascular Health Study	EA	Baseline	1989-90	339	70.6 (3.8)	210 (62%)	368 (74.15, 826.65)	311 (92%)	237 (70%)	147 (43%)	103 (30%)	27 (8%)	39 (12%)	Imatron C-150	27
	AA	Baseline	1989-90, 1992-93	104	70.9 (4.0)	59 (57%)	124.85 (7.15, 499.02)	88 (85%)	55 (53%)	70 (67%)	20 (19%)	21 (21%)	18 (17%)	Imatron C-150	
Diabetes Heart Study	EA	Baseline	1998-2006	667	60.9 (9.5)	416 (62%)	102.5 (12.0, 673.0)	326 (91%)	208 (58%)	544 (81.56 %)	290 (43.67%)	527 (79%)	110 (17%)	General Electric CT system (CTi, LightSpeed QXi&Pro16 system, GE Medical Systems, Waukesha, WI)	28
Jackson Health Study	AA	Baseline	2005-08	1355	59.4 (10.9)	448 (33%)	0.0 (0.0, 65.62)	619 (46%)	283 (21%)	801 (59%)	233 (17%)	178 (13%)	402 (30%)	Lightspeed 16 Pro (GE Healthcare, Milwaukee, WI)	29
Family Heart Study	EA	Baseline	2002-04	2370	55.6 (13.1)	1388 (59%)	0.5 (0.0, 78.0)	1304 (55%)	540 (23%)	855 (36%)	652 (28%)	258 (11%)	234 (10%)	GE LightSpeed Plus, Siemens Volume Zoom, Philips/Marconi MX 8000	5, 30
	AA	Baseline	2002-04	537	52.4 (10.7)	363 (68%)	0 (0.00,28 .00)	266 (50%)	78 (15%)	385 (72%)	117 (22%)	120 (23%)	118 (22%)	GE LightSpeed Plus	
GeneSTAR	EA	Baseline or Follow up	2001, 2003-07, 2008-12	488	49.4 (10.3)	255 (52%)	0 (0, 29.7)	205 (42%)	78 (16%)	187 (38%)	172 (35%)	42 (9%)	83 (17%)	2001-07: Siemens Volume Zoom MDCT; 2008-12: Siemens SOMATOM	31

	AA	Baseline or Follow up	2001, 2003-07, 2008-12	290	50.0 (10.3)	194 (67%)	0 (0, 2.6)	94 (32%)	25 (9%)	163 (56%)	74 (26%)	49 (17%)	81 (28%)	Definition Flash dual-source 256 MDCT 2001-07: Siemens Volume Zoom multidetector row computed tomograph; 2008-12: Siemens SOMATOM Definition Flash dual-source 256 multi-detector scanner	
NELSON	AA	Baseline	2003-06	905	62.0 (6.0)	6 (1%)	97.41 (1.97, 654.47)	706 (78%)	452 (50%)	N/A	N/A	N/A	N/A	Mx8000 IDT, Somatom Sensation 16, Brilliance 16P	32, 33
Old Order Amish	EA	Baseline	2001-06	1075	56.6 (13.1)	591 (55%)	2.40 (0, 144.86)	568 (53%)	304 (26%)	198 (18%)	453 (42%)	45 (4%)	84 (8%)	Imatron C-150	8
FHS	EA	Follow up	2005	3184	52.2 (11.6)	1543(48 %)	0 (0, 45.26)	1204 (38%)	598 (19%)	905 (28%)	722 (23%)	183 (6%)	413 (13%)	LightSpeed Ultra (GE)	9, 10
BioImage	EA	Baseline	2008-09	4182	69.1 (6.0)	2357 (56%)	57.0 (0.0, 274.0)	2983 (71%)	1783 (43%)	2944 (70%)	2713 (65%)	686 (16%)	357 (9%)	Philips Brilliance 64-slice MDCT (Philips Healthcare, Andover, Massachusetts)	11, 34
	AA	Baseline	2008-09	843	67.9 (5.7)	501 (59%)	2.0 (0.0, 77.5)	443 (53%)	179 (21%)	708 (84%)	545 (65%)	250 (30%)	80 (9%)		
MESA	EA	Baseline	2000-02	2526	62.7 (10.2)	1320 (52%)	9.35 (0,6452. 61)	1438 (57%)	780 (31%)	975 (39%)	707 (28%)	151(6%)	287 (11%)	Wake Forest: Light Speed Plus, Light Speed QX/I; Columbia,	30, 35
	AA	Baseline	2000-02	1611	62.3 (10.1)	868 (54%)	0 (0,5599. 43)	721 (45%)	311 (19%)	959(59.53%)	367 (23%)	279 (17%)	194 (13%)	Northwestern, UCLA: Imatron C-150; Johns Hopkins and	

Minnesota: Volume Zoom															
GENOA	EA	Follow up	2000 - 04	1033	58.0 (10.1)	615 (59%)	10.67 (0.0, 141.86)	680 (66%)	298 (29%)	727 (70%)	370 (36%)	136 (13%)	87 (18%)	Imatron C-150	29, 36
	AA	Follow up	2009 - 11	354	67.3 (8.4)	257 (73%)	11.09 (0.0, 196.27)	213 (60%)	114 (32%)	293 (83%)	163 (46%)	120 (34%)	27 (19%)	Lightspeed 16 Pro (GE Healthcare, Milwaukee, WI)	
AGES Reykjavik	EA	Baseline	2002-04	2286	76.2 (5.5)	1457 (63.8%)	(22.59,604.76)	1948 (85%)	1377 (60%)	1763 (77%)	NA	236 (10%)	296 (13%)	Siemens Somatom Sensation 4 MDCT (Siemens Medical Solutions, Malvern, PA)	37
Rotterdam Study	EA	Follow up	1997-99	785	71.8 (5.9)	353 (45%)	146.47 (20.50, 539.09)	734 (94%)	443 (56%)	502 (64%)	311 (40%)	97 (12%)	165 (21%)	C-150 Imatron EBCT (GE-Imatron Inc., South San Francisco, CA)	38
Young Finns Study	EA	Follow up	2007	499	41.7 (2.6)	284 (57%)	0.00 (0.00, 0.00)	96 (19%)	13 (3%)	136 (27%)	76 (15%)	7 (1%)	83 (17%)	GE Discovery VCT 64-slice CT/PET (GE Healthcare, Milwaukee, WI) (Turku), Philips Brilliance 64-slice CT (Philips Medical Systems, Best, Netherlands) (Tampere), Siemens Somatom Sensation 16-	39

	slice CT (Siemens Healthcare, Erlangen, Germany) (Kuopio)
--	--

Table S2. Demographics of Participants by Cohort in CIMT Analysis

Cohort	Ethnicity	Years measured		N	Age, y mean(sd)	Women, n (%)	CIMT, mm median (IQR)	Carotid plaque, n (%)	Hypertension, n (%)	Hypercholesterolemia, n (%)	Diabetes mellitus, n (%)	Recent cigarette smoker, n (%)	Device	References
Cardiovascular Health Study	EA	Baseline	1989-90	4040	72.8(5.6)	2276 (56%)	1.015 (0.91, 1.14)	2722 (70%)	2234 (55%)	1033 (26%)	557 (14%)	426 (11%)	B-mode US (model SSA- 270A; Toshiba)	40
	AA	Baseline	1989-90, 1992-93	738	72.8(5.7)	461 (62%)	1.09 (0.97, 1.22)	552 (75%)	536 (73%)	174 (24%)	166 (23%)	122 (17%)		
Diabetes Heart Study	EA	Baseline	1998-2006	647	60.5(9.7)	413 (64%)	0.631 (0.567, 0.706)	NA	527 (81%)	272 (42%)	510 (79%)	113 (18%)	B-mode US (model AU5; Biosound Esaote)	41
Jackson Heart Study	AA	Baseline	2000-04	1979	60.0(10.5)	678 (34%)	0.70 (0.60, 0.82)	NA	1209 (61%)	348 (18%)	285 (14%)	621 (31%)	B-mode US (Hewlett Packard Sonos 4500)	42
Lothian Birth Cohort 1936	EA	Follow up	2007-10	746	72.8(0.8)	346 (48%)	0.85 (0.70, 0.95)	NA	NA	NA	NA	NA	B-mode US	6
SHIP	EA	Baseline	1997-2001	3504	52.8(13.5)	1742 (50%)	0.81 (0.24)	1864 (53%)	1891 (54%)	NA	403 (12%)	985 (28%)	B-Mode US (Diasonics VST Gateway, Santa Clara, California, USA)	7, 43
SHIP-Trend	EA	Baseline	2008-12	3461	52.1(15.3)	1770 (51%)	0.72 (0.24)	1425 (41%)	1646 (48%)	NA	433 (13%)	868 (25%)	B-Mode US (vivid-i, GE Medical Systems, Waukesha, Wisconsin, WI, USA)	7, 43
GeneSTAR	EA	Follow	2009-12	441	50.0(11.3)	236	0.679	NA	147 (33%)	148 (34%)	37 (8%)	74 (17%)	B-mode US	31, 44

	AA	up Follow up	2009-12	251	49.9(11.8)	(54%) 158 (63%)	(0.504, 0.854) 0.721 (0.547, 0.895)	NA	130 (52%)	82 (33%)	43 (17%)	69 (27%)	(model Philips CX50)	
FHS	EA	Baseline	1995-99	3030	58.5(9.7)	1614 (53%)	0.70 (0.63, 0.80)	534 (18%)	1218 (40%)	869 (29%)	297 (10%)	473 (16%)	Toshiba SSH- 140A imaging unit with a 7.0 MHz transducer	9, 10
BioImage	EA	Baseline	2008-09	4378	69.1(6.0)	2449 (56%)	0.73 (0.65, 0.84)	3487 (80%)	3098 (71%)	2845 (65%)	722 (17%)	380 (9%)	Philips iU22 US systems (Philips Healthcare, Bothell, Washington)	11, 34
	AA	Baseline	2008-09	885	68.0(5.7)	528 (60%)	0.75 (0.68, 0.87)	572 (65%)	748 (85%)	566 (64%)	269 (30%)	85 (10%)	Philips iU22 US systems (Philips Healthcare, Bothell, Washington)	
MESA	EA	Baseline	2000-02	2501	62.6(10.3)	1307 (53%)	0.84 (0.47,2. 45)	NA	963 (39%)	698(28%)	150 (6%)	282 (11%)	B-mode US (Logiq 700 ultrasound device (GE Med Systems, Waukesha, WI))	45, 46
	AA	Baseline	2000-02	1573	62.2(10.3)	848 (54%)	0.89(0.4 0,2.39)	NA	928 (59%)	357(23%)	273 (17%)	291 (19%)		
AGES Reykjavik	EA	Baseline	2002-04	2835	73.6(5.5)	1629 (58%)	1.13 (1.02,1. 24)	NA	2288 (81%)	NA	329 (12%)	357 (13%)	B-Mode US	37
ARIC	EA	Baseline	1987-89	8668	54.3(5.7)	4594 (47%)	0.76 (0.65, 0.85)	1628 (19%)	2301 (27%)	2216 (26%)	723 (8%)	2155 (25%)	B-mode US	15
	AA	Baseline	1987-89	2662	53.9(5.9)	1647 (62%)	0.78 (0.69, 0.90)	457 (17%)	1493 (56%)	685 (27%)	531 (20%)	851 (32%)		
Rotterdam Study	EA	Baseline	1990-93	2500	69.5(8.4)	1341 (54%)	0.99 (0.88,	1451 (58%)	1332 (53%)	1493 (60%)	252 (10%)	607 (24%)	7.5 MHz linear array	47

1.12)													transducer (ATL UltraMark IV; Advanced Technology Laboratories, Bethel, Washington)	
ERF	EA	Baseline	2002-05	1050	49.8(14.2)	564 (54%)	0.78 (0.68, 0.94)	624 (59%)	550 (52%)	364 (35%)	72 (7%)	398 (38%)	7.5 MHz linear-array transducer (ATL 77 UltraMark IV; Advanced Technology Laboratories, Bethel, Washington)	48
NEO	EA	Baseline	2008-12	5666	56.0(5.9)	2943 (52%)	0.99 (0.89, 1.10)	NA	1820 (32%)	1669 (30%)	572 (10%)	896 (16%)	7.5-10 MHz linear-array transducer (Art.Lab v2.1, Esaote, Maastricht, The Netherlands)	19
Young Finns Study	EA	Follow up	2001, 2007	1974	37.3(5.3)	1100 (56%)	0.61 (0.56,0. 67)	54 (3%)	350 (18%)	234 (12%)	23 (1%)	352 (18%)	13.0 MHz linear-array transducer, Sequoia 512 ultrasound mainframes (Acuson Inc., Mountain View, CA)	49

Table S3. *APOE* ε2 association with CAC adjusted for blood lipids.

	All N = 20,527			EA N = 15,398			AA N = 5,129		
Adjustment	Beta	SE	P	Beta	SE	P	Beta	SE	P
Base Model	-0.2519	0.0373	1.54 x 10 ⁻¹¹	-0.1792	0.0442	4.94 x 10 ⁻⁵	-0.4346	0.0700	5.36 x 10 ⁻¹⁰
LDL cholesterol	-0.1862	0.0332	2.08 x 10 ⁻⁸	-0.1748	0.0455	1.23 x 10 ⁻⁴	-0.1993	0.0486	4.14 x 10 ⁻⁵
HDL cholesterol	-0.2061	0.0327	2.95 x 10 ⁻¹⁰	-0.1701	0.0447	1.44 x 10 ⁻⁴	-0.2473	0.0479	2.45 x 10 ⁻⁷
Triglycerides	-0.2260	0.0326	4.45 x 10 ⁻¹²	-0.1942	0.0448	1.44 x 10 ⁻⁵	-0.2619	0.0477	3.95 x 10 ⁻⁸
Total cholesterol	-0.1851	0.0329	1.78 x 10 ⁻⁸	-0.1630	0.0450	2.91 x 10 ⁻⁴	-0.2103	0.0481	1.26 x 10 ⁻⁵

For individuals where fasting lipids were available, the association of ε2 with CAC was repeated individually adjusting for each of the major blood lipid measures within each cohort and meta-analyzed in a fixed effects model. β-Coefficients are estimated for natural log transformation of total Agatston CAC score+1. The base model was adjusted for age, sex, and principal components of ancestry.

AA = African ancestry; CAC = coronary artery calcification; EA = European ancestry; HDL = high density lipoprotein; LDL = low-density lipoprotein; SE = standard error

Table S4. CHD-associated variants and association with subclinical atherosclerosis

Chr	Pos	Gene	SNP	Effect Allele	Effect Allele Frequency	OR	P	CAC			CIMT		
								Beta	SE	P	Beta	SE	P
1	55496039	<i>PCSK9</i>	rs11206510	T	0.85	1.08	2.34E-08	-0.0094	0.0273	7.31E-01	-0.0006	0.0014	6.95E-01
1	56965664	<i>PPAP2B</i>	rs9970807	C	0.92	1.13	5.00E-14	0.0700	0.0353	4.75E-02	0.0033	0.0019	8.09E-02
1	109818530	<i>SORT1</i>	rs646776	T	0.75	1.11	9.01E-19	0.0629	0.0250	1.18E-02	0.0030	0.0013	1.84E-02
1	154422067	<i>IL6R</i>	rs4845625	T	0.45	1.05	3.93E-08	0.0655	0.0212	1.97E-03	-0.0006	0.0011	5.83E-01
1	222823529	<i>MIA3</i>	rs17465637	C	0.66	1.08	3.52E-12	0.0732	0.0233	1.67E-03	0.0002	0.0012	8.65E-01
2	19942473	<i>AKO97927</i>	rs16986953	A	0.11	1.09	1.45E-08	0.1468	0.0423	5.15E-04	-0.0009	0.0021	6.60E-01
2	21286057	<i>APOB</i>	rs515135	C	0.79	1.07	3.09E-08	0.0272	0.0276	3.24E-01	0.0038	0.0014	9.12E-03
2	44073881	<i>ABCG5- ABCG8 VAMP5-</i>	rs6544713	T	0.32	1.05	8.88E-07	0.0240	0.0227	2.90E-01	0.0010	0.0012	3.91E-01
2	85809989	<i>VAMP8- GGCX</i>	rs1561198	T	0.46	1.06	6.37E-10	0.0433	0.0208	3.78E-02	0.0005	0.0011	6.16E-01
2	145801461	<i>ZEB2- ACO74093.1</i>	rs2252641	C	0.48	1.03	5.16E-04	0.0317	0.0210	1.32E-01	0.0001	0.0011	9.35E-01
3	138122122	<i>MRAS</i>	rs9818870	T	0.14	1.07	2.21E-06	0.0338	0.0293	2.48E-01	0.0002	0.0015	8.76E-01
4	156635309	<i>GUCY1A3</i>	rs7692387	G	0.81	1.07	7.35E-09	0.0113	0.0265	6.70E-01	0.0014	0.0014	3.08E-01
6	12903957	<i>PHACTR1</i>	rs9349379	G	0.43	1.14	1.81E-42	0.1922	0.0212	1.28E-19	-0.0006	0.0011	6.00E-01
6	39174922	<i>KCNK5</i>	rs10947789	T	0.78	1.05	1.63E-06	-0.0126	0.0242	6.04E-01	-0.0004	0.0013	7.25E-01
6	160863532	<i>SLC22A3- LPAL2-LPA</i>	rs2048327	C	0.35	1.06	2.46E-09	-0.0041	0.0219	8.52E-01	-0.0014	0.0011	2.20E-01
6	161143608	<i>PLG</i>	rs4252120	T	0.74	1.03	3.32E-03	-0.0080	0.0230	7.29E-01	-0.0014	0.0012	2.28E-01
7	19036775	<i>HDAC9</i>	rs2023938	C	0.1	1.06	1.36E-04	0.0948	0.0349	6.58E-03	0.0044	0.0018	1.44E-02
7	107244545	<i>7q22</i>	rs10953541	C	0.78	1.05	1.02E-05	-0.0207	0.0243	3.95E-01	-0.0002	0.0013	8.96E-01
7	129663496	<i>ZC3HC1</i>	rs11556924	C	0.69	1.08	5.34E-11	0.0377	0.0216	8.17E-02	0.0004	0.0011	7.39E-01
8	19813180	<i>LPL</i>	rs264	G	0.85	1.06	1.06E-05	0.0536	0.0300	7.46E-02	0.0024	0.0015	1.13E-01
8	126490972	<i>TRIB1</i>	rs2954029	A	0.55	1.04	2.61E-06	0.0011	0.0218	9.60E-01	0.0018	0.0012	1.32E-01
9	22098574	<i>9p21</i>	rs4977574	G	0.49	1.21	6.35E-98	0.2111	0.0208	2.88E-24	0.0006	0.0011	5.60E-01
9	136154168	<i>ABO</i>	rs579459	C	0.21	1.08	1.14E-10	0.0450	0.0262	8.62E-02	-0.0030	0.0014	2.81E-02

10	30335122	<i>KIAA1462</i>	rs2505083	C	0.40	1.06	1.57E-10	0.0732	0.0211	5.29E-04	0.0033	0.0011	2.43E-03
10	44753867	<i>CXCL12</i>	rs501120	T	0.81	1.08	1.39E-11	0.0672	0.0312	3.13E-02	-0.0032	0.0016	4.91E-02
10	90989109	<i>LIPA</i>	rs11203042	T	0.45	1.04	1.22E-04	0.0136	0.0210	5.16E-01	-0.0012	0.0011	2.56E-01
10	104719096	<i>CYP17A1- CNNM2- NT5C2</i>	rs12413409	G	0.89	1.08	1.07E-07	-0.0296	0.0377	4.31E-01	0.0014	0.0019	4.50E-01
11	103660567	<i>PDGFD</i>	rs974819	T	0.33	1.07	2.44E-10	0.0818	0.0233	4.56E-04	0.0033	0.0012	7.15E-03
11	116648917	<i>ZNF259- APOA5- APOA1</i>	rs964184	G	0.19	1.05	5.60E-05	0.0288	0.0295	3.30E-01	-0.0010	0.0017	5.45E-01
12	90008959	<i>ATP2B1</i>	rs2681472	G	0.20	1.08	6.17E-11	0.0956	0.0279	6.17E-04	-0.0053	0.0015	2.78E-04
12	111884608	<i>SH2B3</i>	rs3184504	T	0.42	1.07	1.03E-09	-0.0062	0.0208	7.67E-01	-0.0015	0.0011	1.50E-01
13	28973621	<i>FLT1</i>	rs9319428	A	0.31	1.04	7.13E-05	0.0081	0.0224	7.17E-01	-0.0029	0.0012	1.38E-02
14	100133942	<i>HHIPL1</i>	rs2895811	C	0.41	1.04	1.86E-05	0.0283	0.0212	1.81E-01	0.0004	0.0011	6.93E-01
15	91416550	<i>FURIN-FES</i>	rs17514846	A	0.44	1.05	3.10E-07	-0.0410	0.0210	5.08E-02	0.0019	0.0011	8.59E-02
17	2126504	<i>SMG6</i>	rs216172	C	0.35	1.05	5.07E-07	0.0461	0.0217	3.36E-02	-0.0031	0.0012	1.05E-02
17	17543722	<i>RAI1-PEMT- RASD1</i>	rs12936587	G	0.61	1.03	8.24E-04	0.0153	0.0219	4.86E-01	0.0014	0.0011	2.11E-01
17	46988597	<i>UBE2Z</i>	rs46522	T	0.51	1.04	1.84E-05	0.0434	0.0208	3.71E-02	0.0001	0.0011	9.27E-01
19	11163601	<i>LDLR</i>	rs1122608	G	0.77	1.08	2.73E-11	0.0188	0.0243	4.39E-01	0.0036	0.0013	7.59E-03
19	45395619	<i>APOE- APOC1</i>	rs2075650	G	0.13	1.07	1.61E-06	0.0822	0.0299	6.06E-03	0.0018	0.0015	2.23E-01
21	35599128	<i>KCNE2</i> (gene desert)	rs9982601	T	0.13	1.12	1.33E-13	0.0763	0.0324	1.84E-02	-0.0003	0.0016	8.71E-01

Variants that have been previously associated with CHD in individuals of European, South Asian, or East Asian ancestry²⁶, effect estimates, and risk allele frequencies are displayed. The CAC and CIMT effects estimates and significance for each CHD risk variant among those of European ancestry are also displayed. β -Coefficients for CAC and CIMT are estimated for natural log transformation of total Agatston CAC score+1 and natural log transformation of CIMT, respectively.

CAC = coronary artery calcification; CHD = coronary heart disease; Chr = chromosome; CIMT = carotid intima media thickness; OR = odds ratio; P = p-value; Pos = genomic position (hg19 build); SE = standard error; SNP = single nucleotide polymorphism

Supplemental References

1. Grove ML, Yu B, Cochran BJ, Haritunians T, Bis JC, Taylor KD, et al. Best practices and joint calling of the HumanExome BeadChip: the CHARGE Consortium. *PLoS One*. 2013;8:e68095.
2. Bowden DW, Cox AJ, Freedman BI, Hugenschmidt CE, Wagenknecht LE, Herrington D, et al. Review of the Diabetes Heart Study (DHS) family of studies: a comprehensively examined sample for genetic and epidemiological studies of type 2 diabetes and its complications. *Rev Diabet Stud*. 2010;7:188-201.
3. Goldstein JL, Crenshaw A, Carey J, Grant GB, Maguire J, Fromer M, et al. zCall: a rare variant caller for array-based genotyping: genetics and population analysis. *Bioinformatics*. 2012;28:2543-2545.
4. Taylor HA, Jr., Wilson JG, Jones DW, Sarpong DF, Srinivasan A, Garrison RJ, et al. Toward resolution of cardiovascular health disparities in African Americans: design and methods of the Jackson Heart Study. *Ethn Dis*. 2005;15:S6-4-17.
5. Higgins M, Province M, Heiss G, Eckfeldt J, Ellison RC, Folsom AR, et al. NHLBI Family Heart Study: objectives and design. *Am J Epidemiol*. 1996;143:1219-1228.
6. Deary IJ, Gow AJ, Pattie A and Starr JM. Cohort profile: the Lothian Birth Cohorts of 1921 and 1936. *Int J Epidemiol*. 2012;41:1576-1584.
7. Volzke H, Alte D, Schmidt CO, Radke D, Lorbeer R, Friedrich N, et al. Cohort profile: the study of health in Pomerania. *Int J Epidemiol*. 2011;40:294-307.
8. Post W, Bielak LF, Ryan KA, Cheng YC, Shen H, Rumberger JA, et al. Determinants of coronary artery and aortic calcification in the Old Order Amish. *Circulation*. 2007;115:717-724.
9. Kannel WB, Feinleib M, McNamara PM, Garrison RJ and Castelli WP. An investigation of coronary heart disease in families. The Framingham offspring study. *Am J Epidemiol*. 1979;110:281-290.
10. Splansky GL, Corey D, Yang Q, Atwood LD, Cupples LA, Benjamin EJ, et al. The Third Generation Cohort of the National Heart, Lung, and Blood Institute's Framingham Heart Study: design, recruitment, and initial examination. *Am J Epidemiol*. 2007;165:1328-1335.
11. Baber U, Mehran R, Sartori S, Schoos MM, Sillesen H, Muntendam P, et al. Prevalence, Impact, and Predictive Value of Detecting Subclinical Coronary and Carotid Atherosclerosis in Asymptomatic Adults: The BioImage Study. *J Am Coll Cardiol*. 2015;65:1065-1074.
12. Natarajan P, Kohli P, Baber U, Nguyen KD, Sartori S, Reilly DF, et al. Association of APOC3 Loss-of-Function Mutations With Plasma Lipids and Subclinical Atherosclerosis: The Multi-Ethnic BioImage Study. *J Am Coll Cardiol*. 2015;66:2053-2055.
13. Bild DE, Bluemke DA, Burke GL, Detrano R, Diez Roux AV, Folsom AR, et al. Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol*. 2002;156:871-881.
14. Daniels PR, Kardina SL, Hanis CL, Brown CA, Hutchinson R, Boerwinkle E, et al. Familial aggregation of hypertension treatment and control in the Genetic Epidemiology Network of Arteriopathy (GENOA) study. *Am J Med*. 2004;116:676-681.
15. The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. The ARIC investigators. *Am J Epidemiol*. 1989;129:687-702.
16. Hofman A, Brusselle GG, Darwish Murad S, van Duijn CM, Franco OH, Goedegebuure A, et al. The Rotterdam Study: 2016 objectives and design update. *Eur J Epidemiol*. 2015;30:661-708.

17. Aulchenko YS, Heutink P, Mackay I, Bertoli-Avella AM, Pullen J, Vaessen N, et al. Linkage disequilibrium in young genetically isolated Dutch population. *Eur J Hum Genet.* 2004;12:527-534.
18. Pardo LM, MacKay I, Oostra B, van Duijn CM and Aulchenko YS. The effect of genetic drift in a young genetically isolated population. *Ann Hum Genet.* 2005;69:288-295.
19. de Mutsert R, den Heijer M, Rabelink TJ, Smit JW, Romijn JA, Jukema JW, et al. The Netherlands Epidemiology of Obesity (NEO) study: study design and data collection. *Eur J Epidemiol.* 2013;28:513-523.
20. Raitakari OT, Juonala M, Ronnemaa T, Keltikangas-Jarvinen L, Rasanen L, Pietikainen M, et al. Cohort profile: the cardiovascular risk in Young Finns Study. *Int J Epidemiol.* 2008;37:1220-1226.
21. Peloso GM, Auer PL, Bis JC, Voorman A, Morrison AC, Stitzel NO, et al. Association of low-frequency and rare coding-sequence variants with blood lipids and coronary heart disease in 56,000 whites and blacks. *Am J Hum Genet.* 2014;94:223-232.
22. Liu X, Jian X and Boerwinkle E. dbNSFP v2.0: a database of human non-synonymous SNVs and their functional predictions and annotations. *Hum Mutat.* 2013;34:E2393-2402.
23. Baigent C, Keech A, Kearney PM, Blackwell L, Buck G, Pollicino C, et al. Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis. *Lancet.* 2005;366:1267-1278.
24. Eichner JE, Dunn ST, Perveen G, Thompson DM, Stewart KE and Stroehla BC. Apolipoprotein E polymorphism and cardiovascular disease: a HuGE review. *Am J Epidemiol.* 2002;155:487-495.
25. Stitzel NO, Stirrups K, Masca NGD, Erdmann J, Ferrario PG, Konig IR, et al. Coding Variation in ANGPTL4, LPL, and SVEP1 and the Risk of Coronary Disease. *N Engl J Med.* 2016;374:1134-1144.
26. CARDIoGRAMplusC4D. A comprehensive 1000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet.* 2015;47:1121-1130.
27. Newman AB, Naydeck B, Sutton-Tyrrell K, Edmundowicz D, Gottdiener J and Kuller LH. Coronary artery calcification in older adults with minimal clinical or subclinical cardiovascular disease. *J Am Geriatr Soc.* 2000;48:256-263.
28. Carr JJ, Register TC, Hsu FC, Lohman K, Lenchik L, Bowden DW, et al. Calcified atherosclerotic plaque and bone mineral density in type 2 diabetes: the diabetes heart study. *Bone.* 2008;42:43-52.
29. Tullos BW, Sung JH, Lee JE, Criqui MH, Mitchell ME and Taylor HA. Ankle-brachial index (ABI), abdominal aortic calcification (AAC), and coronary artery calcification (CAC): the Jackson heart study. *Int J Cardiovasc Imaging.* 2013;29:891-897.
30. Carr JJ, Nelson JC, Wong ND, McNitt-Gray M, Arad Y, Jacobs DR, Jr., et al. Calcified coronary artery plaque measurement with cardiac CT in population-based studies: standardized protocol of Multi-Ethnic Study of Atherosclerosis (MESA) and Coronary Artery Risk Development in Young Adults (CARDIA) study. *Radiology.* 2005;234:35-43.
31. Kral BG, Becker LC, Vaidya D, Yanek LR, Qayyum R, Zimmerman SL, et al. Noncalcified coronary plaque volumes in healthy people with a family history of early onset coronary artery disease. *Circ Cardiovasc Imaging.* 2014;7:446-453.
32. Isgum I, Prokop M, Niemeijer M, Viergever MA and van Ginneken B. Automatic coronary calcium scoring in low-dose chest computed tomography. *IEEE Trans Med Imaging.* 2012;31:2322-2334.

33. van Setten J, Isgum I, Smolonska J, Ripke S, de Jong PA, Oudkerk M, et al. Genome-wide association study of coronary and aortic calcification implicates risk loci for coronary artery disease and myocardial infarction. *Atherosclerosis*. 2013;228:400-405.
34. Sillesen H, Muntendam P, Adourian A, Entrekin R, Garcia M, Falk E, et al. Carotid plaque burden as a measure of subclinical atherosclerosis: comparison with other tests for subclinical arterial disease in the High Risk Plaque BioImage study. *JACC Cardiovasc Imaging*. 2012;5:681-689.
35. Detrano R, Guerci AD, Carr JJ, Bild DE, Burke G, Folsom AR, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med*. 2008;358:1336-1345.
36. Bielak LF, Sheedy PF, 2nd and Peyser PA. Coronary artery calcification measured at electron-beam CT: agreement in dual scan runs and change over time. *Radiology*. 2001;218:224-229.
37. Harris TB, Launer LJ, Eiriksdottir G, Kjartansson O, Jonsson PV, Sigurdsson G, et al. Age, Gene/Environment Susceptibility-Reykjavik Study: multidisciplinary applied phenomics. *Am J Epidemiol*. 2007;165:1076-1087.
38. Elias-Smale SE, Proenca RV, Koller MT, Kavousi M, van Rooij FJ, Hunink MG, et al. Coronary calcium score improves classification of coronary heart disease risk in the elderly: the Rotterdam study. *J Am Coll Cardiol*. 2010;56:1407-1414.
39. Hartiala O, Magnussen CG, Kajander S, Knuuti J, Ukkonen H, Saraste A, et al. Adolescence risk factors are predictive of coronary artery calcification at middle age: the cardiovascular risk in young Finns study. *J Am Coll Cardiol*. 2012;60:1364-1370.
40. O'Leary DH, Polak JF, Wolfson SK, Jr., Bond MG, Bommer W, Sheth S, et al. Use of sonography to evaluate carotid atherosclerosis in the elderly. The Cardiovascular Health Study. CHS Collaborative Research Group. *Stroke*. 1991;22:1155-1163.
41. Lange LA, Bowden DW, Langefeld CD, Wagenknecht LE, Carr JJ, Rich SS, et al. Heritability of carotid artery intima-medial thickness in type 2 diabetes. *Stroke*. 2002;33:1876-1881.
42. Gebreab SY, Riestra P, Khan RJ, Xu R, Musani SK, Tekola-Ayele F, et al. Genetic ancestry is associated with measures of subclinical atherosclerosis in African Americans: the Jackson Heart Study. *Arterioscler Thromb Vasc Biol*. 2015;35:1271-1278.
43. Ludemann J, Piek M, Wood WG, Meyer S, Greiner B, John U, et al. [Methods for quality assurance of medical examination in epidemiological field studies: the "Study of Health in Pomerania" (SHIP)]. *Gesundheitswesen*. 2000;62:234-243.
44. Kral BG, Nyquist P, Vaidya D, Yousem D, Yanek LR, Fishman EK, et al. Relation of subclinical coronary artery atherosclerosis to cerebral white matter disease in healthy subjects from families with early-onset coronary artery disease. *Am J Cardiol*. 2013;112:747-752.
45. Gepner AD, Young R, Delaney JA, Tattersall MC, Blaha MJ, Post WS, et al. Comparison of coronary artery calcium presence, carotid plaque presence, and carotid intima-media thickness for cardiovascular disease prediction in the Multi-Ethnic Study of Atherosclerosis. *Circ Cardiovasc Imaging*. 2015;8.
46. Polak JF, Szklo M, Kronmal RA, Burke GL, Shea S, Zavodni AE, et al. The value of carotid artery plaque and intima-media thickness for incident cardiovascular disease: the multi-ethnic study of atherosclerosis. *J Am Heart Assoc*. 2013;2:e000087.

47. Bots ML, Hoes AW, Koudstaal PJ, Hofman A and Grobbee DE. Common carotid intima-media thickness and risk of stroke and myocardial infarction: the Rotterdam Study. *Circulation*. 1997;96:1432-1437.
48. Sayed-Tabatabaei FA, van Rijn MJ, Schut AF, Aulchenko YS, Croes EA, Zillikens MC, et al. Heritability of the function and structure of the arterial wall: findings of the Erasmus Rucphen Family (ERF) study. *Stroke*. 2005;36:2351-2356.
49. Bis JC, Kavousi M, Franceschini N, Isaacs A, Abecasis GR, Schminke U, et al. Meta-analysis of genome-wide association studies from the CHARGE consortium. *Nat Genet*. 2011;43:940-947.